

A SOLUTION OF ONE PROBLEM OF J. LINDENSTRAUSS AND H.P. ROSENTHAL ON SUBSPACE HOMOGENEOUS AND QUOTIENT HOMOGENEOUS BANACH SPACES WITH APPLICATION TO THE APPROXIMATION PROBLEM AND TO THE SCHROEDER - BERNSTEIN PROBLEM

EUGENE V. TOKAREV

Dedicated to the memory of S. Banach.

ABSTRACT. In article is constructed a wide couple of pairwise non isomorphic separable superreflexive Banach spaces E that are subspace homogeneous. Their conjugates are quotient homogeneous. None of this couple neither isomorphic to its Cartesian square nor has the approximation property. At the same time, any such E is isomorphic to $E \oplus E \oplus W$ for a some Banach space W and, hence, solves the Schroeder - Bernstein problem.

1. INTRODUCTION

Notions of subspace- and of quotient-homogeneous Banach spaces were introduced by J. Lindenstrauss and H.P. Rosenthal [1] (definitions will be given later). They showed that l_2 and c_0 are subspace-homogeneous, that l_2 and l_1 are quotient-homogeneous and interested in following questions:

1. *Whether there exists separable subspace-homogeneous Banach spaces that are different from l_2 and c_0 ?*
2. *Whether there exists separable quotient homogeneous Banach spaces that are different from l_2 and l_1 ?*
3. *Whether there exists non separable subspace- (or quotient-) homogeneous Banach spaces?*

In the article it will be shown that answers on the first and on the second question are affirmative. Any separable superreflexive Banach space X may be isomorphically (even $(1 + \varepsilon)$ -isomorphically) embedded in a subspace-homogeneous separable superreflexive Banach space E_X of the same type and cotype as X . Of course, $(E_X)^*$ is quotient homogeneous (by superreflexivity of E_X and by duality). Methods that are presented below give no a chance to establish the existence of nonreflexive B-convex spaces with such properties.

The third question is more complicated. A succeed in resolving this problem depends on set-theoretical hypothesis. It may be proved that if there exists a regular cardinal $\varkappa$ with a property: $\varkappa = \varkappa^*$, where $\varkappa^* = \sup\{2^\tau : \tau < \varkappa\}$, then there exists a Banach space of dimension $\varkappa$ which is subspace-homogeneous. Recall

1991 *Mathematics Subject Classification.* Primary 46B10; Secondary 46A20, 46B07, 46B20, 46B28.

Key words and phrases. Banach spaces, Duality, Quotient-closed classes of Banach spaces.

that the existence of a such cardinal $\aleph$ cannot be proved in ZFC. Author expect to present this result in other paper.

Unexpectedly it turned out, that spaces E_X mentioned above posses additional peculiar property. Namely, if there exists a real number $p \neq 2$ such that l_p is finite representable in X , then E_X does not enjoy the approximation property.

This gives an independent solution of the *approximation problem*, that was solved be P. Enflo [2].

If X is not isomorphic to a Hilbert space, then E_X and $E_X \oplus E_X$ are not isomorphic (notice, this fact gives an independent solution of *Banach's problem on squares*, that was solved firstly in [3] and [4]).

Any E_X is universal for all separable spaces that are finitely representable in E_X . This result is related to a *problem on existing of separable envelopes*: any E_X is an "almost envelope": if Z is finitely representable in E_X then, for any $\varepsilon > 0$, Z is $(1 + \varepsilon)$ -isomorphic to a subspace of E_X .

Any E_X has a complement in any Banach space Y that contains E_X and is finitely representable in E_X . In particular, $E_X \oplus E_X$ is isomorphic to a complemented subspace of E_X . Since E_X obviously is a complemented subspace of $E_X \oplus E_X$, a pair $\langle E_X, E_X \oplus E_X \rangle$ solves the *Schroeder-Bernstein problem* that in other way was solved by W.T. Gowers [5].

2. DEFINITIONS AND NOTATIONS

Definition 1. A Banach space X is said to be subspace homogeneous if for any pair of its subspaces Y and Z both of equal codimension (finite or infinite) such that there exists an isomorphism $u : Y \rightarrow Z$ there exists an isomorphical automorphism $U : X \rightarrow X$ such that its restriction $U|_Y = u$.

Definition 2. A Banach space X is said to be quotient homogeneous if for any pair of its quotients X/Y and X/Z such that both Y and Z are of equal (finite or infinite) dimension and there exists an isomorphism $u : X/Y \rightarrow X/Z$, there exists also an automorphism $U : X \rightarrow X$ such that $u \circ h_Y = h_Z \circ U$, where $h_Y : X \rightarrow X/Y$ and $h_Z : X \rightarrow X/Z$ are standard quotient maps.

Definition 3. Let X, Y are Banach spaces. X is finitely representable in Y (in symbols: $X <_f Y$) if for each $\varepsilon > 0$ and for every finite dimensional subspace A of X there exists a subspace B of Y and an isomorphism $u : A \rightarrow B$ such that $\|u\| \|u^{-1}\| \leq 1 + \varepsilon$.

Spaces X and Y are said to be finitely equivalent if $X <_f Y$ and $Y <_f X$.

Any Banach space X generates a class

$$X^f = \{Y \in \mathcal{B} : X \sim_f Y\}$$

Let $\mathcal{K}$ be a class of Banach spaces; X be a Banach space. A notion $X <_f \mathcal{K}$ (or, equivalently, $X \in \mathcal{K}^{<f}$) means that $X <_f Y$ for every $Y \in \mathcal{K}$.

For any two Banach spaces X, Y their *Banach-Mazur distance* is given by

$$d(X, Y) = \inf\{\|u\| \|u^{-1}\| : u : X \rightarrow Y\},$$

where u runs all isomorphisms between X and Y and is assumed, as usual, that $\inf \emptyset = \infty$.

It is well known that $\log d(X, Y)$ defines a metric on each class of isomorphic Banach spaces. A set $\mathfrak{M}_n$ of all n -dimensional Banach spaces, equipped with this metric, is a compact metric space that is called *the Minkowski compact* $\mathfrak{M}_n$.

A disjoint union $\cup\{\mathfrak{M}_n : n < \infty\} = \mathfrak{M}$ is a separable metric space, which is called *the Minkowski space*.

Consider a Banach space X . Let $H(X)$ be a set of all its *different* finite dimensional subspaces (*isometric finite dimensional subspaces of X in $H(X)$ are identified*). Thus, $H(X)$ may be regarded as a subset of $\mathfrak{M}$, equipped with the restriction of the metric topology of $\mathfrak{M}$.

Of course, $H(X)$ need not to be a closed subset of $\mathfrak{M}$. Its closure in $\mathfrak{M}$ will be denoted $\overline{H(X)}$. From definitions follows that $X <_f Y$ if and only if $\overline{H(X)} \subseteq \overline{H(Y)}$. Certainly, spaces X and Y are *finitely equivalent* ($X \sim_f Y$) if and only if $\overline{H(X)} = \overline{H(Y)}$.

Thus, there is a one to one correspondence between classes of finite equivalence $X^f = \{Y \in \mathcal{B} : X \sim_f Y\}$ and closed subsets of $\mathfrak{M}$ of kind $\overline{H(X)}$.

Indeed, all spaces Y from X^f have the same set $\overline{H(X)}$. This set, uniquely determined by X (or, equivalently, by X^f), will be denoted by $\mathfrak{M}(X^f)$ and will be referred as to *the Minkowski's base of the class X^f* .

Using this correspondence, it may be defined a set $f(\mathcal{B})$ of all different classes of finite equivalence, assuming (to exclude contradictions with the set theory) that members of $f(\mathcal{B})$ are sets $\mathfrak{M}(X^f)$. For simplicity it may be says that members of $f(\mathcal{B})$ are classes X^f itself.

Clearly, $f(\mathcal{B})$ is partially ordered by the relation $\mathfrak{M}(X^f) \subseteq \mathfrak{M}(Y^f)$, which may be replaced by the relation $X^f <_f Y^f$ of the same meaning. The minimal (with respect to this order) element of $f(\mathcal{B})$ is the class $(l_2)^f$ (the Dvoretzki theorem); the maximal one - the class $(l_\infty)^f$ (an easy consequence of the Hahn-Banach theorem). Other l_p 's are used in the classifications of Banach spaces, which was proposed by L. Schwartz [6].

For a Banach space X its l_p -spectrum $S(X)$ is given by

$$S(X) = \{p \in [0, \infty] : l_p <_f X\}.$$

Certainly, if $X \sim_f Y$ then $S(X) = S(Y)$. Thus, the l_p -spectrum $S(X)$ may be regarded as a property of the whole class X^f . So, notations like $S(X^f)$ are of obvious meaning.

Let X be a Banach space. It is called:

- *c-convex*, if $\infty \notin S(X)$;
- *B-convex*, if $1 \notin S(X)$;
- *Finite universal*, if $\infty \in S(X)$.
- *Superreflexive*, if every space of the class X^f is reflexive.

Equivalently, X is superreflexive if any $Y <_f X$ is reflexive. Clearly, any superreflexive Banach space is *B-convex*.

3. QUOTIENT CLOSED AND DIVISIBLE CLASSES OF FINITE EQUIVALENCE

In this section it will be shown how to enlarge a Minkowski's base $\mathfrak{M}(X^f)$ of a certain *B-convex* class X^f to obtain a set $\mathfrak{N}$, which will be a Minkowski's base $\mathfrak{M}(W^f)$ for some class W^f that holds the *B-convexity* (and, also, the l_p -spectrum) of the corresponding class X^f and will be quotient closed.

Definition 4. A class X^f (and its Minkowski's base $\mathfrak{M}(X^f)$) is said to be *divisible* if some (equivalently, any) space $Z \in X^f$ is finitely representable in any its subspace of finite codimension.

Definition 5. Let $\{X_i : i \in I\}$ be a collection of Banach spaces. A space

$$l_2(X_i, I) = \left(\sum \oplus \{X_i : i \in I\} \right)_2$$

is a Banach space of all families $\{x_i \in X_i : i \in I\} = \mathfrak{x}$, with a finite norm

$$\|\mathfrak{x}\|_2 = \sup\{(\sum \{\|x_i\|_{X_i}^2 : i \in I_0\})^{1/2} : I_0 \subset I; \text{card}(I_0) < \infty\}.$$

Example 1. Any Banach space X may be isometrically embedded into a space

$$l_2(X) = (\sum_{i < \infty} \oplus X_i)_2,$$

where all X_i 's are isometric to X . Immediately, $l_2(X)$ generates a divisible class $D_2(X^f) = (l_2(X))^f$ which with the same l_p -spectrum as X^f and is superreflexive if and only if X^f is superreflexive too.

Remark 1. The procedure $D_2 : X^f \rightarrow (l_2(X))^f$ may be regarded as a closure operator on the partially ordered set $f(\mathcal{B})$. Indeed, it is

- Monotone, i.e., $X^f <_f D_2(X^f)$;
- Idempotent, i.e., $D_2(X^f) = D_2(D_2(X^f))$;
- Preserve the order: $X^f <_f Y^f \implies D_2(X^f) <_f D_2(Y^f)$.

It is of interest that extreme points of $f(\mathcal{B})$ are stable under this procedure:

$$D_2((l_2)^f) = (l_2)^f; \quad D_2((c_0)^f) = (c_0)^f.$$

To distinguish between general divisible classes and classes of type $D_2(X^f)$, the last ones will be referred as to *2-divisible classes*.

Definition 6. A class X^f (and its Minkowski's base $\mathfrak{M}(X^f)$) is said to be *quotient closed* if for any $A \in \mathfrak{M}(X^f)$ and its subspace $B \hookrightarrow A$ the quotient A/B belongs to $\mathfrak{M}(X^f)$.

Let $K \subseteq \mathfrak{M}$ be a class of finite dimensional Banach spaces (recall, that isometric spaces are identified). Define operations H , Q and $*$ that transform a class K in another class - $H(K)$; $Q(K)$ or $(K)^*$ respectively. Namely, let

$$\begin{aligned} H(K) &= \{A \in \mathfrak{M} : A \hookrightarrow B; \quad B \in K\} \\ Q(K) &= \{A \in \mathfrak{M} : A = B/F; \quad F \hookrightarrow B; \quad B \in K\} \\ (K)^* &= \{A^* \in \mathfrak{M} : A \in K\} \end{aligned}$$

In words, $H(K)$ consists of all subspaces of spaces from K ; $Q(K)$ contains all quotient spaces of spaces of K ; $(K)^*$ contains all conjugates of spaces of K .

The following theorem lists properties of these operations. In iteration of the operations parentheses may be omitted.

Thus, $K^{**} \stackrel{\text{def}}{=} ((K)^*)^*$; $HH(K) \stackrel{\text{def}}{=} H(H(K))$ and so on.

Theorem 1. Any set K of finite dimensional Banach spaces has the following properties:

1. $K^{**} = K$; $HH(K) = H(K)$; $QQ(K) = Q(K)$;
2. $K \subset H(K)$; $K \subset Q(K)$;
3. If $K_1 \subset K_2$ then $H(K_1) \subset H(K_2)$ and $Q(K_1) \subset Q(K_2)$;
4. $(H(K))^* = Q(K^*)$; $(Q(K))^* = H(K^*)$;
5. $HQ(HQ(K)) = HQ(K)$; $QH(QH(K)) = QH(K)$.

Proof. 1, 2 and 3 are obvious.

4. If $A \in Q(K)$ then $A = B/E$ for some $B \in K$ and its subspace E . Thus, A^* is isometric to a subspace of B^* . Hence, $A^* \in H(B^*)$, i.e., $A^* \in H(K^*)$. Since A is arbitrary, $(Q(K))^* \subseteq H(K^*)$. Analogously, if $B \in K$ and $A \in H(B)$ then A^* may be identified with a quotient $B^*/A^\perp$, where $A^\perp$ is the annihilator of A in B^* :

$$A^\perp = \{f \in B^* : (\forall a \in A) \quad (f(a) = 0)\}.$$

Hence, $A^* \in (Q(K^*))^*$, and thus $(H(K))^* \subseteq Q(K^*)$.

From the other hand,

$$\begin{aligned} H(K^*) &= (H(K^*))^{**} \subseteq (Q(K^{**}))^* = (Q(K))^* ; \\ Q(K^*) &= (Q(K^*))^{**} \subseteq (H(K^{**}))^* = (H(K))^* . \end{aligned}$$

5. Let $A \in HQ(K)$. Then A is isometric to a subspace of some quotient space E/F , where $E \in K$; $F \hookrightarrow E$. If B is a subspace of A then $(A/B)^* = (E/F)^*/B^\perp$, i.e. $(Q(HQ(K)))^* \subseteq Q(Q(K)^*)$. Because of

$$\begin{aligned} Q(HQ(K)) &= (Q(HQ(K)))^{**} \subseteq (Q(Q(K)^*))^* \\ &\subseteq H(Q(K)^{**}) = HQ(K), \end{aligned}$$

we have

$$H(Q(H(Q(K)))) \subseteq H(H(Q(K))) = HQ(K).$$

Analogously, if $A \in QH(K)$, then A is isometric to a quotient space F/E , where $F \in H(B)$ for some $B \in K$ and $E \hookrightarrow B$. If $W \in H(A)$, i.e., if $W \in H(F/E)$, then $W^* = (F/E)^*/W^\perp$ and $(F/E)^*$ is isometric to a subspace $E^\perp$ of $F^* \in (H(B))^*$. Thus, $(H(QH(K)))^* \subseteq H((H(K))^*)$ and

$$\begin{aligned} H(QH(K)) &= (H(QH(K)))^{**} \subseteq (H((H(K))^*))^* \\ &\subseteq Q(H(K)^{**}) = QH(K). \end{aligned}$$

Hence,

$$Q(H(Q(HQ(K)))) \subseteq Q(Q(H(K))) = QH(K).$$

Converse inclusion follows from 2. ■

Consider for a given 2-divisible class X^f its Minkowski's base $\mathfrak{M}(X^f)$ and enlarge it by addition to $\mathfrak{M}(X^f)$ of all quotient spaces of spaces from $\mathfrak{M}(X^f)$ and all their subspaces. In the formal language, consider a set $H(Q(\mathfrak{M}(X^f)))$.

For any class W^f the set $\mathfrak{N} = \mathfrak{M}(W^f)$ has following properties:

- (C) $\mathfrak{N}$ is a closed subset of the Minkowski's space $\mathfrak{M}$;
- (H) If $A \in \mathfrak{N}$ and $B \in H(A)$ then $B \in \mathfrak{N}$;
- (A₀) For any $A, B \in \mathfrak{N}$ there exists $C \in \mathfrak{N}$ such that $A \in H(C)$ and $B \in H(C)$.

Theorem 2. Let $\mathfrak{N}$ be a set of finite dimensional Banach spaces; $\mathfrak{N} \subset \mathfrak{M}$. If $\mathfrak{N}$ has properties (C), (H) and (A₀) then there exists a class X^f such that $\mathfrak{N} = \mathfrak{M}(X^f)$.

Proof. Since $(\mathbf{A}_0)$, all spaces from $\mathfrak{N}$ may be directed in an inductive isometric system. Its direct limit W generates a class W^f with $\mathfrak{M}(W^f) \supseteq \mathfrak{N}$. As it follows from the property $(\mathbf{C})$, $\mathfrak{M}(W^f)$ contains no spaces besides that of $\mathfrak{N}$. Hence, $\mathfrak{M}(W^f) = \mathfrak{N}$.

Another proof may be given by using an ultraproduct of all spaces from $\mathfrak{N}$ by the ultrafilter, coordinated with a partial ordering on $\mathfrak{N}$, which is generated by the property $(\mathbf{A}_0)$. ■

Let X be a B -convex Banach space.

Let $X = l_2(Y)$ (and, hence, $X^f = D_2(Y^f)$). Consider the Minkowski's base $\mathfrak{M}(X^f)$ and its enlargement $H(Q(\mathfrak{M}(X^f))) = HQ\mathfrak{M}(X^f)$.

Theorem 3. *There exists a Banach space W such that $HQ\mathfrak{M}(X^f) = \mathfrak{M}(W^f)$.*

Proof. Obviously, $HQ\mathfrak{M}(X^f)$ has properties $(\mathbf{H})$ and $(\mathbf{C})$.

Since $\mathfrak{M}(X^f)$ is 2-divisible, then for any $A, B \in \mathfrak{M}(X^f)$ a space $A \oplus_2 B$ belongs to $\mathfrak{M}(X^f)$ and, hence, to $HQ\mathfrak{M}(X^f)$.

If $A, B \in Q\mathfrak{M}(X^f)$ then $A = F/F_1$; $B = E/E_1$ for some $E, F \in Q\mathfrak{M}(X^f)$.

Since $F/F_1 \oplus_2 E$ is isometric to a space $(F \oplus_2 E)/F'_1$, where $F'_1 = \{(f, 0) \in F \oplus E : f \in F_1\}$, $F/F_1 \oplus_2 E$ belongs to $Q\mathfrak{M}(X^f)$.

Clearly,

$$F/F_1 \oplus_2 E/E_1 = (F/F_1 \oplus_2 E)/E'_1,$$

where

$$E'_1 = \{(o, e) \in F/F_1 \oplus_2 E : e \in E_1\},$$

and, hence, belongs to $Q\mathfrak{M}(X^f)$ too.

If $A, B \in HQ\mathfrak{M}(X^f)$ then $A \hookrightarrow E$, $B \hookrightarrow F$ for some $E, F \in Q\mathfrak{M}(X^f)$. Since $E \oplus_2 F \in Q\mathfrak{M}(X^f)$, obviously $A \oplus_2 B \in HQ\mathfrak{M}(X^f)$.

Thus, $HQ\mathfrak{M}(X^f)$ has a property $(\mathbf{A}_0)$. A desired result follows from the preceding theorem. ■

Definition 7. *Let X be a Banach space, which generates a class X^f of finite equivalence. A class $** (X^f)$ is defined to be a such class W^f that $\mathfrak{M}(W^f) = HQ\mathfrak{M}(D_2 Y^f)$*

Clearly, W^f is quotient closed. Obviously, $X^f <_f W^f$. It will be said that W^f is a result of a procedure $**$ that acts on $f(\mathcal{B})$.

Remark 2. *The procedure $** : X^f \rightarrow ** (X^f) = W^f$ may be regarded as a closure operator on the partially ordered set $f(\mathcal{B})$. Indeed, it is*

- *Monotone, i.e., $X^f <_f ** (X^f)$;*
- *Idempotent, i.e., $X^f = ** (** (X^f))$;*
- *Preserve the order: $X^f <_f Y^f \implies ** (X^f) <_f ** (Y^f)$.*

It is of interest that extreme points of $f(\mathcal{B})$ are stable under this procedure:

$$** ((l_2)^f) = (l_2)^f ; \quad ** ((c_0)^f) = (c_0)^f .$$

Theorem 4. *For any Banach space X a class $** (X^f)$ is 2-divisible.*

Proof. Let $\mathfrak{N} = \mathfrak{M}(\star\star(X^f))$. Since for any pair $A, B \in \mathfrak{N}$ their l_2 -sum belongs to $\mathfrak{N}$, then by an induction, $(\sum_{i \in I} \oplus A_i)_2 \in \mathfrak{N}$ for any finite subset $\{A_i : i \in I\} \subset \mathfrak{N}$.

Hence, any infinite direct l_2 -sum $(\sum_{i \in I} \oplus A_i)_2$ is finite representable in $\star\star(X^f)$. Let $\{A_i : i < \infty\} \subset \mathfrak{N}$ is dense in $\mathfrak{N}$. Let $Y_1 = (\sum_{i < \infty} \oplus A_i)_2$; $Y_{n+1} = Y_n \oplus_2 Y_1$; $Y_\infty = \lim_{\rightarrow} Y_n$. Clearly, $Y_\infty = l_2(Y)$ and belongs to $\star\star(X^f)$. ■

Let $\star : f(\mathcal{B}) \rightarrow f(\mathcal{B})$ be one else procedure that will be given by following steps.

Let $X \in \mathcal{B}$; $Y^f = D_2(X^f)$. Let $\mathfrak{Y}_0$ be a countable dense subset of $\mathfrak{M}(Y^f)$; $\mathfrak{Y}_0 = (Y_n)_{n < \infty}$. Consider a space $Z = (\sum_{n < \infty} \oplus Y_n)_2$ and its conjugate Z^* . Z^* generates a class $(Z^*)^f$ which will be regarded as a result of acting of a procedure $\star : X^f \rightarrow (Z^*)^f$. Since $(Z^*)^f$ is 2-divisible, iterations of the procedure $\star$ are given by following steps: Let $\mathfrak{Z}_0$ is a countable dense subset of $\mathfrak{M}((Z^*)^f)$; $\mathfrak{Z}_0 = (Z_n)_{n < \infty}$. Consider a space $W = (\sum_{n < \infty} \oplus Z_n)_2$ and its conjugate W^* . Clearly, W^* generates a class

$$(W^*)^f = \star(Z^*)^f = \star\star(X^f).$$

Theorem 5. *For any Banach space X classes $\star\star(X^f)$ and $\star\star(X^f)$ are identical.*

Proof. From the construction follows that $H(Z^*) = (QH(l_2(X)))^*$ and that

$$\begin{aligned} H(W^*) &= (QH(Z^*))^* = (Q(QH(l_2(X))))^* \\ &= H(QH(l_2(X)))^{**} = HQH(l_2(X)). \end{aligned}$$

Hence,

$$\mathfrak{M}((W^*)^f) = HQ\mathfrak{M}(Y^f) = HQ\mathfrak{M}(D_2(X^f)).$$

■

Theorem 6. *Let X be a Banach space, which generates a class of finite equivalence X^f . If X is B -convex, then the procedure $\star\star$ maps X^f to a class $\star\star(X^f)$ with the same lower and upper bounds of its l_p -spectrum as X^f . If X is superreflexive then $\star\star(X^f)$ is superreflexive too.*

Proof. Obviously, if $p \in S(X)$ and $p \leq 2$ then the whole interval $[p, 2]$ (that may consist of one point) belongs to $S(X)$ and, hence, to $S(\star\star(X^f))$. If $p \in S(X)$ and $p > 2$ then, by duality, $p/(p-1) \in S(\star(X^f))$; hence $[p/(p-1), 2] \subset S(\star(X^f))$ and, thus $[2, p] \subset S(X) \subset S(\star\star(X^f))$. If $p \notin S(X)$ then $p/(p-1) \notin S(\star(X^f))$ by its construction and, hence, $p \notin S(\star\star(X^f))$ by the same reason. Hence, if

$$p_X = \inf S(X); \quad q_X = \sup S(X)$$

then

$$S(\star\star(X^f)) = [p_X, q_X] = [\inf S(X), \sup S(X)].$$

The second assertion of the theorem is also obvious. ■

Remark 3. *If X is not B -convex, then*

$$\star(X^f) = \star\star(X^f) = \star\star(X^f) = (c_0)^f.$$

4. SPACES OF ALMOST UNIVERSAL DISPOSITION

Recall a definition, which is due to V.I. Gurarii [7].

Definition 8. Let X be a Banach space; $\mathcal{K}$ be a class of Banach spaces. X is said to be a space of almost universal disposition with respect to $\mathcal{K}$ if for any pair of spaces A, B of $\mathcal{K}$ such that A is a subspace of B ($A \hookrightarrow B$), every $\varepsilon > 0$ and every isomorphic embedding $i : A \rightarrow X$ there exists an isomorphic embedding $\hat{i} : B \rightarrow X$, which extends i (i.e., $\hat{i}|_A = i$) and such, then

$$\|\hat{i}\| \|\hat{i}^{-1}\| \leq (1 + \varepsilon) \|i\| \|i^{-1}\|.$$

In the construction of the classical Gurarii's space of almost universal disposition with respect to $\mathfrak{M}$ the main role plays a property of $\mathfrak{M}$, which may be called the *isomorphic amalgamation property*.

Definition 9. Let $X \in \mathcal{B}$ generates a class X^f with a Minkowski's base $\mathfrak{M}(X^f)$. It will be said that $\mathfrak{M}(X^f)$ (and the whole class X^f) has the *isomorphic amalgamation property* if for any fifth $\langle A, B_1, B_2, i_1, i_2 \rangle$, where $A, B_1, B_2 \in \mathfrak{M}(X^f)$; $i_1 : A \rightarrow B_1$ and $i_2 : A \rightarrow B_2$ are isomorphic embeddings, there exists a triple $\langle j_1, j_2, F \rangle$, where $F \in \mathfrak{M}(X^f)$ and $j_1 : B_1 \rightarrow F$; $j_2 : B_2 \rightarrow F$ are isometric embedding, such that $j_1 \circ i_1 = j_2 \circ i_2$.

Theorem 7. Any 2-divisible quotient closed class X^f has the isomorphic amalgamation property.

Proof. Let $A, B_1, B_2 \in \mathfrak{M}(X^f)$; $i_1 : A \rightarrow B_1$ and $i_2 : A \rightarrow B_2$ are isomorphic embeddings. Since X^f is 2-divisible, $C = B_1 \oplus_2 B_2 \in \mathfrak{M}(X^f)$. Consider a subspace H of C that is formed by elements of kind $(i_1 a, -i_2 a)$, where a runs A :

$$H = \{(i_1 a, -i_2 a) : a \in A\}.$$

Consider a quotient $W = C/H$. Since X^f is quotient closed, $W \in \mathfrak{M}(X^f)$. Let $h : C \rightarrow W$ be a standard quotient map. Let $j_1 : B_1 \rightarrow W$ and $j_2 : B_2 \rightarrow W$ are given by:

$$\begin{aligned} j_1(b_1) &= h(b_1, 0) \quad \text{for } b_1 \in B_1; \\ j_2(b_2) &= h(0, b_2) \quad \text{for } b_2 \in B_2. \end{aligned}$$

It is clear that j_1 and j_2 are isometric embeddings such that $j_1 \circ i_1 = j_2 \circ i_2$. ■

Remark 4. It may be shown that any quotient closed divisible (not necessary 2-divisible) class X^f enjoys the isomorphic amalgamation property too. For aims of the article the preceding result is satisfactory.

The proof of a following results is almost literally repeats the Gurarii's proof [7] on existence of a space of almost universal disposition with respect to $\mathfrak{M}$. Only changes that need to be made are: a substitution of a set $\mathfrak{M}$ with $\mathfrak{M}(X^f)$ for a given class X^f and using instead the mentioned above isomorphic amalgamation property of a set $\mathfrak{M}$ the same property of a set $\mathfrak{M}(X^f)$. For this reason the proof of it is omitted.

Theorem 8. Any $**$ -closed class X^f contains a separable space E_X of almost universal disposition with respect to a set $\mathfrak{M}(X^f)$. This space is unique up to almost isometry and is almost isotropic (in an equivalent terminology, has an almost

transitive norm: for any two elements $a, b \in E_X$, such that $\|a\| = \|b\|$ and every $\varepsilon > 0$ there exists an automorphism $u = u(a, b, \varepsilon) : E_X \xrightarrow{\text{onto}} E_X$ such that $\|u\| \|u^{-1}\| \leq 1 + \varepsilon$ and $ua = b$). This space is an approximative envelope of a class X^f : for any $\varepsilon > 0$ every separable Banach space which is finitely representable in X^f is $(1 + \varepsilon)$ -isomorphic to a subspace of E_X .

5. SUBSPACE HOMOGENEOUS AND QUOTIENT HOMOGENEOUS BANACH SPACES

Theorem 9. Let $X^f = ** (X^f)$ be a superreflexive class; E_X be a corresponding separable space of almost universal disposition. Then E_X is subspace homogeneous.

Proof. Let Z_1 and Z_2 be subspaces of E_X that are of equal codimension in E_X (i.e. $\text{codim}(Z_1) = \dim(E_X/Z_1) = \dim(E_X/Z_2) = \text{codim}(Z_2)$).

If $\text{codim}(Z_1) < \omega$ (here and below ω denotes the first infinite ordinal) then assertions of the theorem are satisfied independently of a property of E_X to be a space of almost universal disposition. Indeed, for an arbitrary Banach space X any its subspaces of equal codimension, say, Y and Z , are isomorphic and each of them has a complement in X . So, X may be represented either as a direct sum $X = Y \oplus A$ or as $X = Z \oplus B$ where $\dim(A) = \dim(B) < \omega$.

Let $v : A \rightarrow B$ and $u : Y \rightarrow Z$ be isomorphisms. Clearly, $v \oplus u : X \rightarrow X$ be an automorphism of X that extends u .

Assume that Z_1 and Z_2 are isomorphic subspaces of E_X of infinite codimension. Let $u : Z_1 \rightarrow Z_2$ be the corresponding isomorphism.

Let Z_1 be represented as a direct limit (= as a closure of the union) of a chain

$$Z'_1 \hookrightarrow Z'_2 \hookrightarrow \dots \hookrightarrow Z'_n \hookrightarrow \dots \hookrightarrow Z_1;$$

namely, $Z_1 = \overline{\cup\{Z'_n : n < \omega\}}$ and $\cup Z'_n$ is dense in Z_1 . Clearly,

$$uZ'_1 \hookrightarrow uZ'_2 \hookrightarrow \dots \hookrightarrow uZ'_n \hookrightarrow \dots \hookrightarrow uZ_1 = Z_2$$

and the union $\cup uZ'_n$ is dense in Z_2 .

Let $(e_n)_{n < \omega}$ be a sequence of linearly independent elements of E_X of norm one, which linear span is dense in E_X .

Denote a restriction $u|_{Z'_n}$ by u_n and define two sequences (f_n) and (g_n) of elements of E_X and a sequence (v_n) of isomorphisms by an induction.

Let f_1 be an element of $(e_n)_{n < \omega}$ with a minimal number, which does not belongs to Z'_1 . Let $W_1 = \text{span}(Z'_1 \cup \{f_1\})$ (as $\text{span}(A)$ here and below is denoted a closure of the linear span of A). Let $\varepsilon > 0$ and $v_1 : W_1 \rightarrow E_X$ be an extension of u_1 (i.e. $v_1|_{W_1} = u_1$) such that $\|v_1\| \|v_1^{-1}\| \leq (1 + \varepsilon) \|u_1\| \|u_1^{-1}\|$. Let $g_1 = v_1 f_1$; $U_1 = v_1 W_1$.

Let g_2 be an element of $(e_n)_{n < \omega}$ with a minimal number, which does not belongs to $\text{span}(uZ'_1 \cup U_1)$ and $U_2 = \text{span}(uZ'_1 \cup U_1 \cup \{g_2\})$. Since E_X is a space of almost universal disposition, there exists an isomorphism $(v_2)^{-1}$ which extends both $(u_2)^{-1}$ and $(v_1)^{-1}$ and satisfies

$$\|v_2\| \|v_2^{-1}\| \leq (1 + \varepsilon^2) \max\{\|u_2\| \|u_2^{-1}\|; \|v_1\| \|v_1^{-1}\|\}.$$

Let $f_2 = (v_2)^{-1} g_2$. This close the first step of the induction.

Assume that $\{f_1, f_2, \dots, f_n\}$; $\{g_1, g_2, \dots, g_n\}$; $\{v_1, v_2, \dots, v_n\}$; $\{U_1, U_2, \dots, U_n\}$ and $\{W_1, W_2, \dots, W_n\}$ are already chosen. If n is odd, we choose sequentially:

f_{n+1} as an element of $(e_n)_{n < \omega}$ with a minimal number, which does not belongs to Z'_{n+1} ; $W_{n+1} = \text{span}(Z'_{n+1} \cup \{f_{n+1}\})$; $v_{n+1} : W_{n+1} \rightarrow E_X$ be an extension of

u_{n+1} such that

$$\|v_{n+1}\| \|v_{n+1}^{-1}\| \leq (1 + \varepsilon^{n+1}) \|u_{n+1}\| \|u_{n+1}^{-1}\|;$$

$$g_{n+1} = v_{n+1} f_{n+1}; \quad U_{n+1} = v_{n+1} W_{n+1}.$$

If n is even, we choose sequentially:

g_{n+1} as an element of $(e_n)_{n < \omega}$ with a minimal number, which does not belongs to $\text{span}(uZ'_n \cup U_n)$; $U_{n+1} = \text{span}(uZ'_n \cup U_n \cup \{g_{n+1}\})$; $(v_{n+1})^{-1}$ which extends both $(u_{n+1})^{-1}$ and $(v_n)^{-1}$ and satisfies

$$\|v_{n+1}\| \|v_{n+1}^{-1}\| \leq (1 + \varepsilon^2) \max\{\|u_{n+1}\| \|u_{n+1}^{-1}\|; \|v_n\| \|v_n^{-1}\|\};$$

$$f_{n+1} = (v_{n+1})^{-1} g_{n+1}$$

Since E_X is superreflexive, a sequence of isomorphisms (v_n) converges to an automorphism $V : E_X \rightarrow E_X$, which satisfies

$$\|V\| \|V^{-1}\| \leq \prod_{n=1}^{\infty} (1 + \varepsilon^n) \|u\| \|u^{-1}\|.$$

■

Corollary 1. *Let $X^f = ** (X^f)$ be a superreflexive class; E_X be a corresponding separable space of almost universal disposition. Then $(E_X)^*$ is quotient homogeneous.*

Proof. The desired result is a consequence of the superreflexivity and follows by duality. ■

6. THE BANACH PROBLEM ON SQUARES

A classical Banach's problem on squares (whether every Banach space is isomorphic to its Cartesian square?) was solved in [2]. At the same time there appeared another solution [3]; one more counterexample was presented in [8].

Superreflexive spaces of almost universal disposition form a wide class of spaces that are not isomorphic to their Cartesian squares, which is different from listed above.

Theorem 10. *Let $**X^f$ be a superreflexive class; E_X be a corresponding separable space of almost universal disposition. Then either E_X is isomorphic to the Hilbert space l_2 or spaces E_X and $E_X \oplus E_X$ are not isomorphic.*

Proof. Let the symbol $Y \approx Z$ means that Y and Z are isomorphic.

Assume that $E_X \approx E_X \oplus E_X$. Then $E_X^* \approx E_X^* \oplus E_X^*$. Since E_X^* is not isomorphic to l_2 , there exists a subspace W^* of E_X^* such that $E_X^* \oplus W^*$ and W^* are not isomorphic. Clearly, W^* may be identified with $(E_X/Y)^*$ for some $Y \hookrightarrow E_X$. Certainly, $E_X \oplus (E_X/Y)$ and E_X/Y are not isomorphic.

Let $u : E_X \rightarrow E_X \oplus E_X$ be an isomorphism. Then $uY \hookrightarrow E_X \oplus E_X$ and $uY \approx Y$.

Consider two isomorphic embeddings, say, j_1 and j_2 of Y into E_X .

Let $j_1 : Y \hookrightarrow E_X$ be the identical embedding. Let $j_2 = u \circ j_1$.

Let $v : j_1 Y \rightarrow j_2 Y$ be an isomorphism. Then $(v^{-1} \circ j_2) Y \hookrightarrow E_X$. However the isomorphism v cannot be extended to any automorphism of E because of absence of isomorphisms between $E_X \oplus (E_X/Y)$ and E_X/Y . But this contradicts with the property of E_X to be subspace homogeneous. ■

Remark 5. If a Banach space X is not isomorphic to any space Y that generates a divisible class Y^f , then any space $Z \in X^f$ is not isomorphic to its Cartesian square. First example of such X was the space $F = \left(\sum \oplus l_{p_i}^{(n_i)} \right)_2$, where $p_i \searrow 2$; $n_i = \dim \left(l_{p_i}^{(n_i)} \right) \rightarrow \infty$; $(p_i - 2) \log n_i \rightarrow \infty$, that was constructed by T. Figiel [8].

7. THE SCHROEDER-BERNSTEIN PROBLEM

The *Schroeder-Bernstein problem*, that was induced by the classical paper [9] and received its name thanks to a certain analogy may be formulated as follows:

Let X, Y, Z be Banach spaces. Is there true that a condition $X \approx X \oplus Y \oplus Z$ implies that $X \approx X \oplus Y$?

Recall that in [9] was shown that if $X \approx X \oplus X$ and $Y \approx Y \oplus Y$ then $X \approx X \oplus Y \oplus Z$ implies that $X \approx X \oplus Y$.

This problem was (in negative) solved by W.T. Gowers (see [5]).

An another solution of this problem based on the property of spaces of kind E_X , which needs additional definitions.

Definition 10. (Cf.[10]). Let X be a Banach space; Y - its subspace. Y is said to be a reflecting subspace of X (symbolically: $Y \prec_u X$) if for every $\varepsilon > 0$ and every finite dimensional subspace $A \hookrightarrow X$ there exist an isomorphic embedding $u : A \rightarrow Y$ such that $\|u\| \|u^{-1}\| \leq 1 + \varepsilon$ and $u|_{A \cap Y} = Id_{A \cap Y}$.

As was shown in [10], if $Y \prec_u X$ then Y^{**} is an image of a norm one projection $P : X^{**} \rightarrow Y^{**}$ (under canonical embedding of Y^{**} into X^{**}).

Definition 11. (Cf [11]). A Banach space E is said to be existentially closed in a class X^f if for any isometric embedding $i : E \rightarrow Z$ into an arbitrary space $Z \in X^f$ its image iE is a reflecting subspace of Z : $iY \prec_u Z$.

A class of all spaces E that are existentially closed in X^f is denoted by $\mathcal{E}(X^f)$. In [11] was shown that for any Banach space X the class $\mathcal{E}(X^f)$ is nonempty; moreover, any $Y \prec_f X^f$ may be isometrically embedded into some $E \in \mathcal{E}(X^f)$ of the dimension $\dim(E) = \max\{\dim(Y), \omega\}$.

Theorem 11. For any class $** (X^f)$ the corresponding space E_X of almost universal disposition is existentially closed in $** (X^f)$.

Proof. Let $E_X \hookrightarrow Z \in ** (X^f)$. Let $A \hookrightarrow Z$. Denote $E_X \cap A = B$. Since $E_X \sim_f Z$, for any $\varepsilon > 0$ there exists an isomorphic embedding $u : A \rightarrow E_X$ such that $\|u\| \|u^{-1}\| \leq 1 + \varepsilon/4$. Let $F = uA$; $G = u(E_X \cap A) = uB$. Since $F \subset E_X$ and $G \subset E_X$, a restriction $u|_B = v$ may be extended to an isomorphism $V : F \rightarrow E_X$ such that $\|V\| \|V^{-1}\| \leq 1 + \varepsilon/4$. Clearly, VF is $(1 + \varepsilon)$ -isomorphic to A and $VF \cap A = B$. Because of A is arbitrary, $E_X \prec_u Z$ and, hence, $E_X \in \mathcal{E}(** (X^f))$. ■

Corollary 2. If $** (X^f)$ is superreflexive, then E_X is a norm one complemented subspace of any space Z from the class $** (X^f)$ that it contain.

Proof. If $E_X \hookrightarrow Z \in ** (X^f)$ then $E_X \prec_u Z$ and hence $(E_X)^{**} = E_X$ is a norm one complemented subspace of $Z^{**} = Z$ according to [10]. ■

Corollary 3. Thou $E_X \approx E_X \oplus E_X \oplus Z$ but E_X and $E_X \oplus E_X$ are not isomorphic.

Proof. Because $E_X \oplus_2 E_X$ is finitely equivalent to E_X , it may be isomorphically embedded into E_X . Since the image of E_X in E_X is a complemented subspace, clearly, $E_X \oplus E_X$ is isomorphic to a complemented subspace of E_X , i.e., $E_X \approx E_X \oplus E_X \oplus Z$. In a previous section was proved that E_X and $E_X \oplus E_X$ are not isomorphic. ■

8. THE APPROXIMATION PROBLEM

Definition 12. A Banach space X has the approximation property (shortly: AP) provided for every compact subset $K \subset X$ and every $\varepsilon > 0$ there exists a (bounded linear) operator $u_K : X \rightarrow X$ with a finite dimensional range such that $\|u_K x - x\| \leq \varepsilon$ for all $x \in K$.

If there exists a constant λ , independent on K and ε , such that norms of operators u_K are uniformly bounded by λ : $\|u_K\| \leq \lambda$, then X is said to have the λ -bounded approximation property (λ -BAP).

If X has the λ -BAP for all $\lambda > 1$, then X is said to have the metric approximation property (MAP).

The approximation problem, i.e. a question: "whether every Banach space has the approximation property?" was solved (in negative) by P. Enflo [2]. This solution (and other known counterexamples to the approximation problem) are quite complicated.

This section is devoted to prove that a lot of spaces of kind E_X - i.e. separable spaces of almost universal disposition, that are presented in every quotient closed and divisible class X^f - does not enjoy the approximation property. Unfortunately, using methods cannot give a chance to prove the absence of approximation property in a non reflexive case. It seems that result presented below also is far from finality.

Theorem 12. Let $** (X^f)$ be a superreflexive class such that its l_p -spectrum $S(** (X^f))$ contains a point $p \neq 2$. Let E_X be the corresponding space of almost universal disposition. Then E_X does not have the approximation property.

Proof. Let $p \in S(** (X^f))$ and $p \neq 2$. Then either $p_X = \inf S(** (X^f)) < 2$ or $q_X = \sup S(** (X^f)) > 2$. From [12] follows that if $r = p_X$ or $r = q_X$ then E_X (as any other space from $** (X^f)$) contains a sequence $(Y_n)_{n < \infty}$ of finite dimensional subspaces of unboundedly growing dimension that are uniformly complemented in E_X and are uniformly isomorphic to l_r^k . More precise, $\dim(Y_n) \rightarrow \infty$ and there exists such a constant C that $\sup d(Y_n, l_r^{\dim(Y_n)}) < C$ and that

$$\sup\{\|P_n\| : P_n : E_X \rightarrow Y_n\}$$

for some sequence of projections (P_n) .

Since E_X is a space of almost universal disposition, it may be assumed without loss of generality that the sequence $(Y_n)_{n < \infty}$ is directed in a chain

$$Y_1 \hookrightarrow Y_2 \hookrightarrow \dots \hookrightarrow Y_n \hookrightarrow \dots \hookrightarrow E_X.$$

From [13] follows that $Y = \overline{\cup_n Y_n}$ is an $\mathcal{L}_p$ -space which cannot have the complement in E_X if $r \neq 2$. Indeed, if Y is a complemented subspace of E_X then E_X contains a complemented subspace isomorphic to l_r . At the same time, E_X contains an uncomplemented subspace, isomorphic to l_r . Indeed (we assume that $r \neq 2$), as an universal space for a class of separable spaces from $(** (X^f))^{<f}$, E_X contains a subspace isomorphic to $L_r[0, 1]$ (for simplicity assume that $L_r[0, 1]$ is a subspace

of E_X itself). The last one contains an uncomplemented subspace Z isomorphic to l_r (for $p > 2$ this fact is contained in [14]; for $p < 2$ follows from [15]). If there is a projection from E_X onto Z then its restriction is a projection from $L_r[0, 1]$ onto Z what is impossible. Thus, E_X contains an uncomplemented subspace, isomorphic to l_r . Since E_X is subspace homogeneous, every its subspace, isomorphic to l_r has no complement in E_X .

Assume that E_X has the approximation property. By the superreflexivity, E_X has the metric approximation property. Hence, there exists a sequence $(E^{(n)})$ of finite dimensional subspaces of E_X and a sequence of operators $R_n : E_X \rightarrow E^{(n)}$ with $\sup \|R_n\| \leq \lambda$ for a given $\lambda > 1$, which convergents pointwise to the identical operator Id_{E_X} . It may be assumed that $Y_n \subset E^{(m_n)}$ for a some sequence (m_n) of natural numbers. A restriction P'_n of P_n to $E^{(m_n)}$ is a projection from $E^{(m_n)}$ onto Y_n . A sequence $(R_{m_n} \circ P'_n)$ convergents pointwise to a projection $P : E_X \rightarrow Y = \overline{\bigcup_n Y_n}$. As it was shown, this is impossible. ■

Problem 1. *Whether non reflexive space of kind E_X either is isomorphic (=isometric) to the classical Gurarii space of almost universal disposition or does not enjoy the approximation property?*

9. TSIRELSON LIKE SPACES

A Banach space X is said to has the Tsirelson property if it contains no subspaces isomorphic to the spaces l_p ($1 \leq p < \infty$) and c_0 .

The first example of a Banach space with a such property was constructed by B.S. Tsirelson [16]. Below it will be shown that spaces that enjoy the Tsirelson property may be found among spaces of kind E_X .

Theorem 13. *Let $** (X^f)$ be a superreflexive class such that its l_p -spectrum $S(** (X^f))$ is a single point $p = 2$: $S(** (X^f)) = \{2\}$. Let E_X be the corresponding space of almost universal disposition. If E_X is not of cotype 2 then either its conjugate $(E_X)^*$ has the Tsirelson property or E_X is isomorphic to the Hilbert space.*

Proof. Assume that E_X is not isomorphic to the Hilbert space.

Since E_X is not of cotype 2, according to the Kwapien's result [17] its 2-type constant $T_2(X)$ is infinite. So, for any number N there exists a finite dimensional subspace $X^{(n)}$ with $T_2(X^{(n)}) \geq N$, where, certainly, $n = \dim(X^{(n)}) = n(N)$ depends on N (and tends to infinity with N).

Fix $\varepsilon > 0$; $1 < C < \infty$ and, proceeding by induction, chose sequences $(p_k) \subset \mathbb{R}_+$; $p_k < p_{k+1} < 2$; $(n_k) \subset \mathbb{N}$ and $(N_k) \subset \mathbb{R}$.

Let $N_1 > 2$ be arbitrary chosen. Let n_1 be the least natural number such that there exists a subspace $X^{(n_1)} \hookrightarrow X$ of dimension n_1 for which

$$T_2(X^{(n_1)}) \geq N_1.$$

Let

$$p_1 = 2 - C(\log(n_1))^{-1}.$$

As N_2 may be chosen any number such that

$$(N_2)^{C(\log(n_1))^{-1}} \geq 2N_1.$$

Assume that $N_1, N_2, \dots, N_k; p_1, p_2, \dots, p_{k-1}; n_1, n_2, \dots, n_{k-1}$ and subspaces $X^{(n_1)}, X^{(n_2)}, \dots, X^{(n_{k-1})}$ are chosen. Let n_k be the least natural number such that there exists a subspace $X^{(n_k)} \hookrightarrow X$ of dimension n_k for which

$$T_2(X^{(n_k)}) \geq N_k.$$

Put

$$p_k = 2 - C (\log(n_k))^{-1}.$$

Let N_{k+1} satisfies

$$(N_{k+1})^{C(\log(n_k))^{-1}} \geq 2N_k.$$

It was noted before that L_p for $p < 2$ has an uncomplemented subspace, isomorphic to l_2 . The finite-dimensional version of this fact is:

There exists such a constant c that for every $N > 1$ there exists such $n = n(N)$ that $l_p^{(n)}$ contains a subspace $Y^{(m_n)}$ of dimension m_n such that

$$d(Y^{(m_n)}, l_2^{(m_n)}) \leq c; \quad \|P\| \geq N \text{ for every projection } P : l_p^{(n)} \rightarrow Y^{(m_n)}.$$

From [18] follows that for any $p \in (1, 2)$ every Banach space X , which type p -constant is $T_p(X)$, contains a subspace, which is $(1 + \varepsilon)$ -isomorphic to $l_p^{(m)}$, of dimension $m = m(p, \varepsilon)$ that is estimated as

$$m = m(p, \varepsilon) \geq C(\varepsilon) (T_p(X))^{1/q},$$

where $p^{-1} + q^{-1} = 1$ and $C(\varepsilon)$ depends only on ε .

From the definition of type p -constants follows that for a finite dimensional Banach space X

$$|T_p(X) - T_2(X)| \leq (\dim(X))^{1/p-1/2}.$$

At least, according to the Kwapien's result, for any Banach space X of cotype 2

$$T_p(X) \asymp d(X, l_2^{\dim(X)}).$$

Since our choosing of $\langle N_k, n_k, p_k, X^{(n_k)} \rangle$'s follows:

1. $|T_p(X^{(n_k)}) - T_2(X^{(n_k)})| \leq C$ for all $k < \infty$;
2. $X^{(n_k)}$ contains a subspace that is $(1 + \varepsilon)$ -isomorphic to $l_p^{(m(p_k, \varepsilon))}$.

Let $m_k = m(p_k, \varepsilon)$. Then

$$\begin{aligned} d(l_{p_k}^{n_k}, l_2^{n_k}) &= (m_k)^{1/p_k-1/2} \geq \left(C(\varepsilon) (T_{p_k}(X^{(n_k)}))^{1/q} \right)^{1/p_k-1/2} \\ &\geq C(\varepsilon) (T_2(X^{(n_k)}) - C)^{1/q} \geq C(\varepsilon) (2^k N_1 - C)^{1/q} \end{aligned}$$

and, hence, tends to infinity with k .

So, E_X (as any other Banach space X of cotype 2 and the single-point spectrum $S(X) = \{2\}$) contains a sequence of finite dimensional subspaces (Y^{m_k}) , which are uniformly isomorphic to Euclidean ones and whose projection constants

$$\lambda(Y^{m_k} \hookrightarrow E_X) = \inf\{\|P\| : P : E_X \rightarrow Y^{m_k}; P^2 = P\}$$

tend to infinity.

Certainly, E_X contains a subspace l_2 (since E_X is an approximate envelope and thanks to the Dvoretzki's theorem).

Clearly, this space has no complement in E_X : in a contrary case any sequence of finite dimensional subspaces of E_X that are uniformly isomorphic to Euclidean ones should have uniformly bounded projection constants.

Assume that $(E_X)^*$ contains a subspace Z , which is isomorphic to l_2 . Then, according to [19], Z is a complemented subspace of $(E_X)^*$ since it is of type 2. Hence, by superreflexivity, $(E_X)^{**} = E_X$ contains a complemented subspace Z^* , which is also isomorphic to the Hilbert space. However, as was shown before, it is impossible. ■

Remark 6. *In the same way (i.e. using the presence in certain spaces of kind E_X of an uncomplemented Hilbertian subspace) it may be obtained the following extension of the theorem 12.*

Theorem 14. *Let $** (X^f)$ be a superreflexive class such that $S(** (X^f)) = \{2\}$. Let E_X be the corresponding space of almost universal disposition. If E_X is not of type 2 then it fails to have the approximation property.*

Problem 2. *Whether superreflexive spaces of kind E_X either are isomorphic to the Hilbert space or has no approximation property?*

10. REFERENCES

1. Lindenstrauss J., Rosenthal H.P. *Automorphisms in c_o , l_1 and m* , Israel J. Math. **7** (1969) 227-239
2. Enflo P. *A counterexample to the approximation problem in Banach spaces*, Acta Math. **130** (1973) 309-317
3. Bessaga C., Pełczyński A. *Banach spaces non - isomorphic to their Cartesian squares*, Bull Acad. Pol. Sci. Ser Math. Astron. et Phys. **8** (1960) 77-80
4. Semadeni Z. *Banach spaces non-isomorphic to their Cartesian squares*, Bull Acad. Pol. Sci. Ser. Math. Astron. et Phys. **8** (1960) 81-84
5. Gowers W.T. *Recent results in the theory of infinite-dimensional Banach spaces*, Proc. of Int. Congress Math. Zurich, v. **2** (1994) 931-942
6. Schwartz L. *Geometry and probability in Banach spaces*, Bull. AMS **4:2** (1981) 135-141
7. Gurarii V.I. *Spaces of universal disposition, isotropic spaces and the Mazur problem on rotations in Banach spaces*, Sibirsk. Mat. Journ. (in Russian) **7** (1966) 1002-1013
8. Figiel T. *An example of infinite dimensional reflexive Banach space non-isomorphic to its Cartesian square*, Studia Math. **42:8** (1972) 295 - 306
9. Pełczyński A. *Projections in certain Banach spaces*, Studia Math. **19** (1960) 209-228
10. Stern J. *Ultrapowers and local properties in Banach spaces*, Trans. AMS **240** (1978) 231-252
11. Tokarev E.V. *Injective Banach spaces in the finite equivalence classes* (transl. from Russian), Ukrainian Mathematical Journal **39:6** (1987) 614-619
12. Maurey B., Pisier G. *Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach*, Studia Math. **58** (1976) 45-90
13. Lindenstrauss J., Pełczyński A. *Absolutely summing operators in $\mathcal{L}_p$ -spaces and their applications*, Studia Math. **29** (1968) 275-326

14. Rosenthal H.P. *Projections onto translation-invariant subspaces of $L_p(G)$* , Memoirs AMS **63** (1968) 1-84
15. Bennet G., Dor L.E., Gudman V., Johnson W.B., Newmann C.M. *On un-complemented subspaces of L_p , $1 < p < 2$* , Israel J. Math. **26** (1977) 178-187
16. Tsirelson B.S. *Not every Banach space contains l_p or c_0* (in Russian), Funct. Analysis and its Appl. **8:2** (1974) 57-60
17. Kwapien S. *Isomorphic characterization of inner product spaces by orthogonal series with vector valued coefficients*, Studia Math. **44** (1972) 583-595
18. Pisier G. *On the dimension of the $l_p(n)$ -subspaces of Banach spaces for $1 < p < 2$* , Trans. AMS **276** (1983) 201-211
19. Maurey B. *Quelques problèmes de factorization d'opérateurs linéaires*, Actes du Congress Int. Math. Vancouver, 1974 v. **2** (1975) 75-79

B.E. UKRECOLAN, 33-81, ISKRINSKAYA STR., 61005, KHARKIV-5, UKRAINE
E-mail address: `tokarev@univer.kharkov.ua`